

Interventions to improve treatment adherence in migrant people living with HIV in the Netherlands: feasibility and efficacy

S. Been¹, D. van de Vijver², J. Smit³, N. Bassant¹, K. Pogány³, A. Verbon⁴.

1. Dep. of Internal Medicine and Infectious Diseases, Erasmus University Medical Center, Rotterdam, The Netherlands. s.been@erasmusmc.nl. 2. Dep. of Viroscience, Erasmus University Medical Center, Rotterdam. 3. Dep. of Infectious Diseases, Maasstad Hospital, Rotterdam, The Netherlands. 4. Dep. of Medical Microbiology and Infectious Diseases, Erasmus University Medical Center, Rotterdam, The Netherlands.

Background

Non-adherence to combination Antiretroviral Therapy (cART) is more common in non-Western migrants living with HIV (MLWH) than in Dutch people living with HIV (DPLWH). This was shown to be secondary to psychosocial factors like more depressive symptoms, feelings of HIV-related stigma, low social support, quality of life and disclosure concerns. Non-adherence has been associated with more virological failure and missed clinical appointments. The objective of this study was to evaluate the feasibility and efficacy of 4 different interventions to improve adherence among MLWH.

Methods

All MLWH visiting the HIV-treatment centers in Rotterdam were asked to participate and fill out questionnaires before and after the interventions to collect information about, among others, demographics, social support, HIV-related stigma, treatment adherence, treatment adherence self-efficacy and quality of life. Additionally, clinical characteristics were collected. The 4 interventions were: Directly Administered Antiretroviral Therapy (DAART), medical group consultation, early detection and treatment of depression, and peer support. Uni- and multivariable analyses were conducted on baseline data from patients who used cART >6 months. The Wilcoxon matched-pairs test, paired samples T-test, McNemar test, and McNemar-Bowker test were used to compare data of the 2 questionnaires.







Figure 1. Flow diagram of patient inclusion



Results

352 patients were included (41% of the eligible MLWH) of whom 301 used cART >6 months (figure 1). 49.4% of the patients was self-reported adherent and HIV-RNA was detectable in 11%. Non-adherent patients experienced less social support, treatment adherence self-efficacy and had a lower educational attainment. The second questionnaire was filled out by 235 (78%) patients. Self-reported adherence increased significantly from 49.4% to 57.0% ($p < 0.05$) (Table 1). A marginally significant decrease was shown of 10.2% to 6.8% of patients with a detectable HIV-RNA ($p = 0.12$).

Interventions tested between July 2013 – July 2015





Peer Support Project	Medical group consult
 <ul style="list-style-type: none"> • Trained peers: 20 • Matches with migrant PLWH: 41 	 <ul style="list-style-type: none"> • Participants: 28 in 5 groups consults • ROAD participants and non-participants • Mean grade for the group consult: 8.2 out of 10!
Early screening and treatment of depression	Directly Administered Antiretroviral Therapy (DAART)
 <ul style="list-style-type: none"> • Hospital Anxiety and Depression Scale: 307 • At risk for a depression based on HADS: 106 • Referred to a psychologist: 23 	 <ul style="list-style-type: none"> • Participants: 3 • Most eligible participants refused this intervention

Table 1: Socio-demographic characteristics and psychosocial variables

N=235	Q1	Q2	P - value	
HIV-RNA >50 copies/ml (%)	24 (10.2)	16 (6.8)	0.12 ^d	
Living situation (%)			0.23 ^e	
With family	90 (38.3)	81 (34.5)		
Alone	85 (36.2)	96 (40.9)		
Single parent	45 (19.1)	44 (18.7)		
Other	15 (6.4)	14 (6.0)		
Employment status (%)			0.23 ^e	
Paid employment	93 (39.6)	89 (37.9)		
Unemployed	67 (28.5)	62 (26.4)		
On sick leave	26 (11.1)	25 (10.6)		
Other	49 (20.9)	59 (25.1)		
Alcohol in the past 30 days (%) ^a	131 (55.7)	122 (51.9)	0.37 ^d	
Alcohol use ≥3 days/week (%) ^a	41 (17.4)	35 (14.9)	0.42 ^d	
Drugs in the past 30 days (%) ^a	39 (16.6)	36 (15.3)	0.72 ^d	
Drugs use ≥3 days/week (%)	24 (10.2)	23 (9.8)		
Psychosocial variables (median, IQR)				
Social support ^a	75 (40.6-90.6)	68.8 (43.8-87.5)	0.89 ^b	0.97 ^c
Internalized HIV-related stigma ^a	15 (12-19)	14 (11-18)	<0.01 ^b	<0.01 ^c
Treatment adherence self-efficacy ^a	105.5 (91.8-116.0)	106 (89.8-115)	0.91 ^b	0.93 ^c
Quality of life (physical) ^a	51.3 (40.7-56.2)	50.4 (41.7-55.6)	0.29 ^b	0.36 ^c
Quality of life (mental) ^a	49.0 (38.9-57.3)	48.8 (37.8-56.3)	0.83 ^b	0.84 ^c
Self-reported adherence (%) ^a				
I – Adherent	81 (34.5)	96 (40.9)	<0.10 ^d	
I – Non-adherent	152 (64.7)	135 (57.4)		
II – Adherent	108 (46.0)	121 (51.5)	0.15 ^d	
II – Non-adherent	125 (53.2)	110 (46.8)		
III – Adherent	116 (49.4)	134 (57.0)	<0.05 ^d	
III – Non-adherent	117 (49.8)	97 (41.3)		

IQR, Interquartile Range.

^a Missing values: alcohol or drugs in the past 30 days (1), alcohol use ≥3 days/week (1), social support (5), internalized HIV related stigma (9), treatment adherence self-efficacy (33), physical quality of life (9), mental quality of life (9), self-reported adherence (2) ^bWilcoxon matched-pairs test; ^cPaired samples T-test; ^dMcNemar; ^eMcNemar-Bowker

Conclusion

In MLWH, DAART and medical group consultation were not feasible interventions. Despite the fact that only a minority of patients had a referral for depression or peer support, self-reported adherence and undetectable HIV-RNA in the total group increased. Although the 'Hawthorne effect' cannot be excluded, our data suggest that attention for depressive symptoms and low social support increases treatment adherence in MLWH.