

HIV infection is independently associated with chronic kidney disease and mild glomerular hyperfiltration, particularly in those of African descent

Katherine W Kooij¹, Liffert Vogt², Ferdinand W Wit^{1,3,4}, Marc van der Valk³, Rosan A van Zoest¹, Maria Prins^{3,5}, Peter Reiss^{1,3,4},
on behalf of the AGE_nIV Cohort Study Group

¹Academic Medical Center (AMC), Department of Global Health and Amsterdam Institute for Global Health and Development; ²AMC, Department of Nephrology; ³AMC, Department of Internal Medicine, Division of Infectious Diseases and Center for Infection and Immunity Amsterdam; ⁴HIV Monitoring Foundation; ⁵Public Health Service of Amsterdam, Department of Infectious Diseases, Amsterdam, Netherlands

Background

HIV-infected individuals are at increased risk of chronic kidney disease (CKD).¹ Traditional risk factors as well as those related to HIV infection and exposure to antiretroviral therapy (ART), particularly tenofovir disoproxil fumarate (TDF), may all contribute.²

¹Yombi JC et al, HIV Medicine 2015; ²Yombi JC et al, AIDS 2014

Methods

Study population

HIV-1 infected and uninfected AGE_nIV Cohort Study participants, aged ≥45 yrs.

Aims

To cross-sectionally compare the prevalence of a low estimated glomerular filtration rate (eGFR), albuminuria and proximal renal tubular dysfunction (PRTD) between HIV-infected and uninfected study participants.

To compare longitudinal eGFR decline during a follow-up up to 4 years (baseline, 2 biennial study visits) in HIV-infected individuals on cART.

Definitions

Low eGFR: eGFR below 60 ml/min, calculated using the CKD Epidemiology Collaboration formula

Albuminuria: urine albumin-creatinine ratio >3mg/mmol.

PRTD: urine retinol-binding protein:creatinine ratio >2.93µg/mmol and/or fractional phosphate excretion >20% with plasma phosphate <0.8 mmol/L

Statistical analyses

Multivariable logistic and linear regression analyses: to assess independent associations between HIV-status and a low eGFR, albuminuria and PRTD, adjusting for traditional CKD risk factors (listed below).

Linear mixed effects models to estimate eGFR decline over 4 year follow-up.


Multiple imputation was performed to handle missing baseline covariates.

Baseline characteristics

	HIV-infected, n=596 % or median (IQR)	HIV-uninfected, n=544 % or median (IQR)
age, years	52.7 (48.3 – 59.4)	52.1 (47.9 – 58.1)
male gender	87.9%	84.7%
African descent ¹	14.3%	6.4%
current / past smoking	32.1% / 35.1%	24.8% / 38.9%
chronic HCV infection ²	3.5%	1.1%
diabetes mellitus ³	6.7%	4.8%
hypertension ⁴	50.2%	37.3%
LDL-C, mmol/L / using statin	3.1 (2.5 – 3.7) / 14.1%	3.3 (2.7 – 3.9) / 7.8%
using cART	95.0%	
regimen containing TDF	73.3%	


HCV, hepatitis C virus; LDL-C, low-density lipoprotein cholesterol; cART, combination antiretroviral therapy; TDF, tenofovir disoproxil fumarate. ¹birth country Suriname(Creole)/Netherlands Antilles/sub-Saharan Africa and/or invalid AGE_nIV measurement due to low reflection; ²detectable hepatitis C virus RNA; ³HbA1c ≥48 mmol/mol and/or blood glucose (fasting/non-fasting) ≥11.1/≥7.0 mmol/L and/or use of antidiabetic drugs; ⁴mean systolic/diastolic blood pressure ≥140/≥90 and/or use of antihypertensive drugs.

Baseline eGFR distribution



- Median eGFR **not significantly different** between HIV-infected (92.6 ml/min, 95% CI 78 – 101) and uninfected individuals (91.3 ml/min, 95% CI 84 – 100), p=0.97
- More **extreme low and high eGFR values** in the HIV-infected study group
- Particularly **high eGFR** was observed in HIV-infected individuals of **African descent**

Baseline low eGFR, albuminuria and PRTD



HIV infection independently[#] associated with:


Low eGFR OR_{HIV} 2.1 (95% CI 1.0 – 4.4) **Albuminuria** OR_{HIV} 5.8 (95% CI 3.7 – 9.0) **PRTD** OR_{HIV} 7.1 (95% CI 4.9 – 10.2)

Within the HIV-positive group:

- **cumulative duration of TDF exposure** independently associated with PRTD (OR 1.2 per year, 95% CI 1.1 – 1.3)
- **historic but not current TDF exposure** borderline independently associated with low eGFR (OR 3.3, 95% CI 0.9 – 11.4).

[#]adjusted for age, gender, ethnicity, smoking status, HCV infection, diabetes mellitus, hypertension, LDL-cholesterol, and use of statins

Longitudinal eGFR decline



Treated HIV infection independently[#] associated with 0.89 mL/min greater yearly eGFR decline (95% CI 0.58 to 1.20 mL/min, P<0.001).

HIV infection on stable cART independently[#] associated with 0.57 mL/min greater yearly eGFR decline (95% CI 0.25 to 0.88 mL/min, P<0.001).

[#]adjusted for age, gender, ethnicity, smoking status, HCV infection, diabetes mellitus, hypertension, LDL-cholesterol, and use of statins

Conclusions

In this cohort of middle-aged HIV-positive and HIV-negative individuals, the majority on TDF containing ART, **HIV was independently associated with prevalent low eGFR, albuminuria and proximal renal tubular dysfunction.**

Both **very low and high eGFR values** were more common in HIV-positive individuals, with higher eGFR particularly associated with being of African descent. This could be an expression of **glomerular hyperfiltration**, a condition associated with increased risk for CKD development. This suggests that **HIV-positive individuals of African descent** might be at **increased risk to progress towards CKD** despite currently having high eGFR.

Furthermore, treated HIV infection was independently associated with **greater eGFR decline** during a follow-up period up to 4 years. To a large extent, the high prevalence and progression of CKD may be due to the frequent and long-term use of TDF in this population.

Correspondence and funding

KW Kooij, MD, PhD fellow, k.kooij@amc.uva.nl +31205663349 Amsterdam Institute for Global Health and Development (AIGHD), Pietersbergweg 17, 1105 BM Amsterdam, the Netherlands
Financial support: The Netherlands Organisation for Health Research and Development (ZonMW) grant nr. 300020007, Stichting AIDS Fonds grant nr. 2009063.
Unconditional grants: Gilead Sciences, ViiV Healthcare, Janssen Pharmaceuticals, Merck & Co, Bristol Myers Squibb