

Prevalence of cardiovascular risk factors among HIV-infected and –uninfected women in the reproductive age in rural South Africa



UMC Utrecht

Marissa J Harmsen, MD, MSc¹, Annelot F Schoffelen, MD, PhD¹, Walter Devillé, MD, PhD^{2,3}, Hugo A Tempelman, MD², Andy I Hoepelman, MD, PhD¹, Diederick E Grobbee³, Roos E Barth, MD, PhD¹

¹ Department of Internal Medicine & Infectious Diseases at the University Medical Center in Utrecht, Utrecht, The Netherlands. ² Ndlovu Care Group in Elandsdoorn, Limpopo, South Africa. ³ Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands.



Background:

As the risk factors for developing hypertensive disorders of pregnancy largely overlap with those for cardiovascular diseases (CVD), the increasing prevalence of cardiovascular risk factors in Sub-Saharan Africa might increase the risk of complications in HIV-infected pregnancies.

Aim To determine the prevalence of CVD risk factors among HIV-infected and HIV-uninfected women in the reproductive age

Table 1. Characteristics of HIV-infected and –uninfected participants

	Total population		Age adjusted <i>p</i> -value	HIV-infected population		Age adjusted <i>p</i> -value
	HIV-infected (N=194) ^a	HIV-uninfected (N=299) ^b		On HAART (N=140) ^c	Not on HAART (N=54) ^d	
Demographic factors						
Age (years)	34.1 ± 6.7	28.8 ± 8.2	<0.001	35.8 ± 5.7	29.8 ± 7.1	<0.001
HIV-related factors						
On HAART	140 (72.2)	-		140 (100.0)	-	
NNRTI-based 1 st line	132 (94.3)	-		132 (94.3)	-	
PI-based 2 nd line	8 (5.7)	-		8 (5.7)	-	
Lifestyle						
Current smoker	13 (6.7)	23 (7.7)	0.68	10 (7.1)	3 (5.6)	0.76
Alcohol consumer	41 (21.1)	80 (26.8)	0.16	25 (17.9)	16 (29.6)	0.07
Physical inactive	89 (45.9)	169 (56.5)	<0.05	62 (44.3)	27 (50.0)	0.47
Physical examination						
Systolic blood pressure	107 [98-118]	108 [100-120]	0.08	105 [98-116]	108.3 [100-123]	0.07
Hypertensive ^e	11 (5.7)	29 (9.7)		5 (3.6)	6 (11.1)	
Body mass index kg/m ²	25.6 ± 6.5	25.9 ± 6.5	0.62	25.6 ± 6.5	25.7 ± 6.6	0.94
Overweight >25	87 (44.8)	153 (51.1)		64 (45.7)	23 (42.6)	
Laboratory values						
CD4 count cells/mm ³	-	-	-	528 ± 235	507 ± 241	0.60
Viral load <50 copies/mL	-	-	-	108 (77.1)	6 (11.1)	<0.001
C-reactive protein, mg/l	4.0 [2-10]	3.0 [2-6]	<0.001	6.0 [2.0-10.0]	2.0 [2.0-7.3]	<0.05
Glucose, mmol/l	4.6 [4.3-5.0]	4.4 [4.1-4.9]	0.88	4.7 [4.3-5.0]	4.5 [4.2-4.9]	0.99
HbA1c%	5.6 [5.3-5.8]	5.5 [5.2-5.8]	0.94	5.6 [5.3-5.8]	5.6 [5.2-5.8]	0.99
Total cholesterol, mmol/l	4.3 ± 1.0	4.0 ± 0.9	<0.001	4.4 ± 0.9	3.9 ± 0.9	<0.001
Triglycerides, mmol/l	0.9 [0.7-1.3]	0.7 [0.5-1.0]	<0.001	1.0 [0.7-1.4]	0.8 [0.6-1.2]	<0.05
HDL-C, mmol/l	1.5 ± 0.5	1.3 ± 0.3	<0.001	1.5 [1.3-1.8]	1.2 [1.0-1.5]	<0.001
LDL-C, mmol/l	2.3 ± 0.8	2.3 ± 0.8	0.47	2.4 ± 0.8	2.2 ± 0.8	0.20
Creatinine, mmol/l	12.3 ± 6.5	14.2 ± 8.2	<0.05	11.9 ± 6.2	13.4 ± 7.2	0.19
Microalbumin, mg/l	12.0 [6.0-22.5]	8.0 [5.0-16.0]	<0.05	12.2 [6.0-25.9]	10.8 [6.8-17.9]	0.38
Albumin/creat ratio>3.4	32 (16.5)	32 (10.7)	0.06	28 (20.0)	4 (7.4)	<0.05

Data are no.(%) of participants, mean±SD or median [interquartile range]

Methods

- Cross-sectional, observational study
- Setting: rural clinic in Limpopo, South Africa.
- Three CVD risk prediction models

Results

The prevalence of CVD risk factors was low. Most women had 10-year CVD risk scores below 1%. After age adjustment, no difference in CVD risk scores: - between HIV-infected group and -uninfected group - or; group on HAART and treatment naïve patients.

Conclusions

1. No significant differences in the prevalence of CVD risk factors between the HIV-infected and –uninfected women of this rural South African population.
2. The prevalence of obesity and physical inactivity highlights the burden of the increase in CVD risk factors in this area.
3. Lifestyle changes and preventive cardiovascular risk management are therefore recommended.
4. Routine prenatal care should be standard care for all women of the reproductive age.




Figure 1. Age adjusted CVD risk scores in HIV-infected vs. –uninfected population. FHS-10, Framingham 10-year heart risk, DAD, Data collection on Adverse Effects of Anti-HIV Drugs Study.